

Passport data

| | |
|--|---|
| Passport descriptors | |
| Prime name | RIESLING WEISS |
| Color of berry skin | BLANC |
| Variety number VIVC | 10077 |
| Country or region of origin of the variety | GERMANY |
| Species | VITIS VINIFERA LINNÉ SUBSP. SATIVA (DE CANDOLLE) HEGI |

| Degree of resistance | | | | | | | | | | | | | | | | | | | |
|---|-------|-------|-------|-------|-------|-------|-------|----------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-----|
| Source code | 20011 | 20035 | 20036 | 20041 | 20046 | 20052 | 20053 | 20054 | 20058 | 20059 | 20064 | 20065 | 20070 | 20071 | 20072 | 20075 | 20076 | 20081 | |
| Accession name | | | | | | | | RIESLING WEISS | | | | | | | | | | | |
| Alternaria rot (leaf) (<i>Alternaria alternata</i> (Fr.) Keissl) | | | | | | | | 7 | | | | | | | | | | | |
| Anthraco se (leaf) (<i>Elsinoë ampelina</i> Shear) | | | | | | | | | 7 | | | | | | | | | | |
| Anthraco se (bunch) (<i>Elsinoë ampelina</i> Shear) | | | | | | | | | | | | | | | | | | | |
| Anthraco se (leaf/shoot) (<i>Elsinoë ampelina</i> Shear) | 1 | | | | | | | 7 | | 1 | | | 7 | 7 | | | | | |
| Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | | | | | | | | | | | 4 | | | | | | | |
| Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | | | | | | | | | | | | | | | | | | |
| Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | | | | | | | | | | | | | | | | | | |
| Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | | | | | | | | | | | 9 | | | | | | | |
| Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | | | | | | | | | | | | | | | | | | |
| Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | | | | | | | | | | | | | | | | | | |
| Botrytis bunch rot (leaf) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | | | | | | | | | | | | | | | | | | | |
| Botrytis bunch rot (bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | 1 4 | | | | | 1 4 | 1 4 | | | | | | | | | | | | 1 4 |
| Botrytis bunch rot (leaf/bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | | | | | | | | | | | | | | | | | | | |
| Brenner (leaf) (<i>Pseudopeziza tracheiphila</i> Müll.-Thurg.) | 1 | | | | | | | | | | | | | | | | | | 3 |
| Cercospora (leaf) | | | | | | | | 7 | | | | | 7 | | 3 | | | | |
| Crown gall (shoot) (<i>Agrobacterium tumefaciens</i> (Smith & Townsend) Conn) | | | | | | | | | | | | | | | | | | | |
| Downy mildew (leaf) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | | 1 | 4 | | | | | | | | | | | | | | | | |
| Downy mildew (bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | | | | | | | | | | | | | | | | | | | |
| Downy mildew (leaf/bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | | | | | | | | | | | | | | | | | | | |
| Excoriose (leaf/shoot) (<i>Phomopsis viticola</i> (Sacc.) Sacc.) | 3 | | | | | | | | | | 5 | | | | | | | | |
| Isariopsis leaf blight (leaf) (<i>Isariopsis clavispora</i> (Berk. & Cooke) Sacc.) | | | | | | | | 7 | | | | | | | | | | | |

