

Passport data

| | |
|--|---|
| Passport descriptors | |
| Prime name | TINTO CAO |
| Color of berry skin | NOIR |
| Variety number VIVC | 12500 |
| Country or region of origin of the variety | PORTUGAL |
| Species | VITIS VINIFERA LINNÉ SUBSP. SATIVA (DE CANDOLLE) HEGI |

| Degree of resistance | | |
|---|-------|-------|
| Source code | 20071 | 20072 |
| Accession name | | |
| Alternaria rot (leaf) (<i>Alternaria alternata</i> (Fr.) Keissl) | | |
| Anthraco (leaf) (<i>Elsinoë ampelina</i> Shear) | | |
| Anthraco (bunch) (<i>Elsinoë ampelina</i> Shear) | | |
| Anthraco (leaf/shoot) (<i>Elsinoë ampelina</i> Shear) | 7 | |
| Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | |
| Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | |
| Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | |
| Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | |
| Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | |
| Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | |
| Botrytis bunch rot (leaf) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | | |
| Botrytis bunch rot (bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | | |
| Botrytis bunch rot (leaf/bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | | |
| Brenner (leaf) (<i>Pseudopeziza tracheiphila</i> Müll.-Thurg.) | | |
| Cercospora (leaf) | | |
| Crown gall (shoot) (<i>Agrobacterium tumefaciens</i> (Smith & Townsend) Conn) | | |

| | | |
|---|--|---|
| Downy mildew (leaf) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | | |
| Downy mildew (bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | | |
| Downy mildew (leaf/bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | | |
| Excoriöse (leaf/shoot) (<i>Phomopsis viticola</i> (Sacc.) Sacc.) | | |
| Isariopsis leaf blight (leaf) (<i>Isariopsis clavispora</i> (Berk. & Cooke) Sacc.) | | |
| Pierce's Disease (<i>Xylella fastidiosa</i> Wells. et al.: grapevine) | | |
| Powdery mildew (leaf) (<i>Erysiphe necator</i> Schwein.) | | |
| Powdery mildew (bunch) (<i>Erysiphe necator</i> Schwein.) | | |
| Powdery mildew (leaf/bunch) (<i>Erysiphe necator</i> Schwein.) | | 2 |
| Ripe rot (leaf) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk) | | |
| Ripe rot (bunch) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk) | | |
| Ripe rot (leaf/bunch) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk) | | |
| White rot (leaf) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow) | | |
| White rot (bunch) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow) | | |
| White rot (leaf/bunch) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow) | | |