

Passport data

Passport descriptors	
Prime name	CHANCELLOR
Color of berry skin	NOIR
Variety number VIVC	2446
Country or region of origin of the variety	FRANCE
Species	VITIS INTERSPECIFIC CROSSING

Degree of resistance							
Source code	3 25	20011	20022	20039	20040	20065	20080
Accession name	SEIBEL 7053	SEIBEL 7053	SEIBEL 7053		SEIBEL 7053		
Alternaria rot (leaf) (<i>Alternaria alternata</i> (Fr.) Keissl)							
Anthraco se (leaf) (<i>Elsinoë ampelina</i> Shear)							
Anthraco se (bunch) (<i>Elsinoë ampelina</i> Shear)							
Anthraco se (leaf/shoot) (<i>Elsinoë ampelina</i> Shear)		5					
Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.)						8	
Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.)							
Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.)				8			
Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.)						7	
Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.)							
Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.)							
Botrytis bunch rot (leaf) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.))							
Botrytis bunch rot (bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.))		5					
Botrytis bunch rot (leaf/bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.))							
Brenner (leaf) (<i>Pseudopeziza tracheiphila</i> Müll.-Thurg.)		5					
Cercospora (leaf)							

Crown gall (shoot) (<i>Agrobacterium tumefaciens</i> (Smith & Townsend) Conn)							
Downy mildew (leaf) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni)	6						
Downy mildew (bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni)	5						
Downy mildew (leaf/bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni)	7		7	3			
Excoriöse (leaf/shoot) (<i>Phomopsis viticola</i> (Sacc.) Sacc.)							
Isariopsis leaf blight (leaf) (<i>Isariopsis clavispora</i> (Berk. & Cooke) Sacc.)							
Pierce's Disease (<i>Xylella fastidiosa</i> Wells. et al.: grapevine)							
Powdery mildew (leaf) (<i>Erysiphe necator</i> Schwein.)					4		3
Powdery mildew (bunch) (<i>Erysiphe necator</i> Schwein.)					2		3
Powdery mildew (leaf/bunch) (<i>Erysiphe necator</i> Schwein.)			7	3			
Ripe rot (leaf) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)							
Ripe rot (bunch) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)							
Ripe rot (leaf/bunch) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)							
White rot (leaf) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)							
White rot (bunch) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)							
White rot (leaf/bunch) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)							