

Passport data

| | |
|--|------------------------------|
| Passport descriptors | |
| Prime name | COUDERC 136- 04 |
| Color of berry skin | NOIR |
| Variety number VIVC | 3005 |
| Country or region of origin of the variety | FRANCE |
| Species | VITIS INTERSPECIFIC CROSSING |

| | |
|---|---|
| Degree of resistance | |
| Source code | 3 |
| Accession name | |
| Alternaria rot (leaf) (<i>Alternaria alternata</i> (Fr.) Keissl) | |
| Anthraco nose (leaf) (<i>Elsinoë ampelina</i> Shear) | |
| Anthraco nose (bunch) (<i>Elsinoë ampelina</i> Shear) | |
| Anthraco nose (leaf/shoot) (<i>Elsinoë ampelina</i> Shear) | |
| Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>eu vitis</i> Luttr.) | |
| Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>eu vitis</i> Luttr.) | |
| Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>eu vitis</i> Luttr.) | |
| Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | |
| Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | |
| Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | |
| Botrytis bunch rot (leaf) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | |
| Botrytis bunch rot (bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | |
| Botrytis bunch rot (leaf/bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | |
| Brenner (leaf) (<i>Pseudopeziza tracheiphila</i> Müll.-Thurg.) | |
| Cercospora (leaf) | |
| Crown gall (shoot) (<i>Agrobacterium tumefaciens</i> (Smith & Townsend) Conn) | |
| Downy mildew (leaf) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | |
| Downy mildew (bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | |
| Downy mildew (leaf/bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) | 5 |
| Excoriose (leaf/shoot) (<i>Phomopsis viticola</i> (Sacc.) Sacc.) | |
| Isariopsis leaf blight (leaf) (<i>Isariopsis clavispora</i> (Berk. & Cooke) Sacc.) | |

| | |
|--|--|
| Pierce's Disease (<i>Xylella fastidiosa</i> Wells. et al.: grapevine) | |
| Powdery mildew (leaf) (<i>Erysiphe necator</i> Schwein.) | |
| Powdery mildew (bunch) (<i>Erysiphe necator</i> Schwein.) | |
| Powdery mildew (leaf/bunch) (<i>Erysiphe necator</i> Schwein.) | |
| Ripe rot (leaf) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk) | |
| Ripe rot (bunch) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk) | |
| Ripe rot (leaf/bunch) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk) | |
| White rot (leaf) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow) | |
| White rot (bunch) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow) | |
| White rot (leaf/bunch) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow) | |