

## Passport data

|  |   |
|--|---|
| Passport descriptors                       |   |
| Prime name                                 | DOBRICIC  |
| Color of berry skin                        | NOIR  |
| Variety number VIVC                        | 3608  |
| Country or region of origin of the variety | CROATIA   |
| Species                                    | VITIS VINIFERA LINNÉ SUBSP. SATIVA (DE CANDOLLE) HEGI |

|   |          |
|---|----------|
| <b>Degree of resistance</b>   |          |
| Source code   | 20068    |
| Accession name  | DOBRITCH |
| <b>Alternaria rot</b> (leaf)<br>( <i>Alternaria alternata</i> (Fr.) Keissl)                                       |          |
| <b>Anthraxnose</b> (leaf)<br>( <i>Elsinoë ampelina</i> Shear)   |          |
| <b>Anthraxnose</b> (bunch)<br>( <i>Elsinoë ampelina</i> Shear)  |          |
| <b>Anthraxnose</b> (leaf/shoot)<br>( <i>Elsinoë ampelina</i> Shear)   |          |
| <b>Black rot</b> (leaf)<br>( <i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.)          |          |
| <b>Black rot</b> (bunch)<br>( <i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.)         |          |
| <b>Black rot</b> (leaf/bunch)<br>( <i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.)    |          |
| <b>Black rot</b> (leaf)<br>( <i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.)       |          |
| <b>Black rot</b> (bunch)<br>( <i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.)      |          |
| <b>Black rot</b> (leaf/bunch)<br>( <i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) |          |
| <b>Botrytis bunch rot</b> (leaf)<br>( <i>Botrytis cinerea</i> (Botrytis cinerea Pers.))                           |          |
| <b>Botrytis bunch rot</b> (bunch)<br>( <i>Botrytis cinerea</i> (Botrytis cinerea Pers.))                          |          |
| <b>Botrytis bunch rot</b> (leaf/bunch)<br>( <i>Botrytis cinerea</i> (Botrytis cinerea Pers.))                     |          |
| <b>Brenner</b> (leaf)<br>( <i>Pseudopeziza tracheiphila</i> Müll.-Thurg.)   |          |
| <b>Cercospora</b> (leaf)  |          |
| <b>Crown gall</b> (shoot)<br>( <i>Agrobacterium tumefaciens</i> (Smith & Townsend) Conn)                          |          |

|   |   |
|---|---|
| <b>Downy mildew</b> (leaf)<br>( <i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni)       |   |
| <b>Downy mildew</b> (bunch)<br>( <i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni)      |   |
| <b>Downy mildew</b> (leaf/bunch)<br>( <i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni) |   |
| <b>Excoriöse</b> (leaf/shoot)<br>( <i>Phomopsis viticola</i> (Sacc.) Sacc.)                             |   |
| <b>Isariopsis leaf blight</b> (leaf)<br>( <i>Isariopsis clavispora</i> (Berk. & Cooke) Sacc.)           |   |
| <b>Pierce's Disease</b><br>( <i>Xylella fastidiosa</i> Wells. et al.: grapevine)                        |   |
| <b>Powdery mildew</b> (leaf)<br>( <i>Erysiphe necator</i> Schwein.)                                     | 2 |
| <b>Powdery mildew</b> (bunch)<br>( <i>Erysiphe necator</i> Schwein.)                                    | 1 |
| <b>Powdery mildew</b> (leaf/bunch)<br>( <i>Erysiphe necator</i> Schwein.)                               |   |
| <b>Ripe rot</b> (leaf)<br>( <i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)                |   |
| <b>Ripe rot</b> (bunch)<br>( <i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)               |   |
| <b>Ripe rot</b> (leaf/bunch)<br>( <i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)          |   |
| <b>White rot</b> (leaf)<br>( <i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)                        |   |
| <b>White rot</b> (bunch)<br>( <i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)                       |   |
| <b>White rot</b> (leaf/bunch)<br>( <i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)                  |   |