

Passport data

| | |
|--|---|
| Passport descriptors | |
| Prime name | KOENIGIN DER WEINGAERTEN |
| Color of berry skin | BLANC |
| Variety number VIVC | 6350 |
| Country or region of origin of the variety | HUNGARY |
| Species | VITIS VINIFERA LINNÉ SUBSP. SATIVA (DE CANDOLLE) HEGI |

| Degree of resistance | | |
|---|-------|-------|
| Source code | 20011 | 20058 |
| Accession name | | |
| Alternaria rot (leaf) (<i>Alternaria alternata</i> (Fr.) Keissl) | | |
| Anthraxnose (leaf) (<i>Elsinoë ampelina</i> Shear) | | 7 |
| Anthraxnose (bunch) (<i>Elsinoë ampelina</i> Shear) | | |
| Anthraxnose (leaf/shoot) (<i>Elsinoë ampelina</i> Shear) | | |
| Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | |
| Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | |
| Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>euvitis</i> Luttr.) | | |
| Black rot (leaf) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | |
| Black rot (bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | |
| Black rot (leaf/bunch) (<i>Guignardia bidwellii</i> (Ellis) Viala & Ravaz f. <i>muscadinii</i> Luttr.) | | |
| Botrytis bunch rot (leaf) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | | |
| Botrytis bunch rot (bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | 3 | |
| Botrytis bunch rot (leaf/bunch) (<i>Botrytis cinerea</i> (Botrytis cinerea Pers.)) | | |
| Brenner (leaf) (<i>Pseudopeziza tracheiphila</i> Müll.-Thurg.) | | |
| Cercospora (leaf) | | |
| Crown gall (shoot) (<i>Agrobacterium tumefaciens</i> (Smith & Townsend) Conn) | | |

| | | |
|--|--|--|
| <p>Downy mildew (leaf) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni)</p> | | |
| <p>Downy mildew (bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni)</p> | | |
| <p>Downy mildew (leaf/bunch) (<i>Plasmopara viticola</i> (Berk. & M.A. Curtis) Berl. & De Toni)</p> | | |
| <p>Excoriöse (leaf/shoot) (<i>Phomopsis viticola</i> (Sacc.) Sacc.)</p> | | |
| <p>Isariopsis leaf blight (leaf) (<i>Isariopsis clavispora</i> (Berk. & Cooke) Sacc.)</p> | | |
| <p>Pierce's Disease (<i>Xylella fastidiosa</i> Wells. et al.: grapevine)</p> | | |
| <p>Powdery mildew (leaf) (<i>Erysiphe necator</i> Schwein.)</p> | | |
| <p>Powdery mildew (bunch) (<i>Erysiphe necator</i> Schwein.)</p> | | |
| <p>Powdery mildew (leaf/bunch) (<i>Erysiphe necator</i> Schwein.)</p> | | |
| <p>Ripe rot (leaf) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)</p> | | |
| <p>Ripe rot (bunch) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)</p> | | |
| <p>Ripe rot (leaf/bunch) (<i>Glomerella cingulata</i> (Stoneman) Spauld. & H. Schrenk)</p> | | |
| <p>White rot (leaf) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)</p> | | |
| <p>White rot (bunch) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)</p> | | |
| <p>White rot (leaf/bunch) (<i>Coniella diplodiella</i> (Speg.) Petrak & Sydow)</p> | | |